

ON $\mathbb{F}_q$ -RATIONAL STRUCTURE OF NILPOTENT CONES IN CLASSICAL LIE ALGEBRAS

SEMRA KAPTANOGLU, BRIAN PARSHALL AND BIN SHU

ABSTRACT. Let $\mathfrak{g}$ be a classical simple Lie algebra over $k = \bar{\mathbb{F}}_q$, defined over a finite field $\mathbb{F}_q$ of characteristic $p > 3$, with correspondence to Frobenius-Lie morphism F . In this paper, we investigate the relation between F -stable restricted modules and closed conical subvarieties defined over $\mathbb{F}_q$ in the null cone $\mathcal{N}(\mathfrak{g})$ of $\mathfrak{g}$. Furthermore, we clearly investigate the $\mathbb{F}_q$ -rational structure for all nilpotent orbits in $\mathfrak{g}$ under the action of adjoint group G when p is good. We also do that for nilpotent orbits in $\text{Lie}(G)$ under the action of G , a simply-connected simple algebraic group over k and defined over $\mathbb{F}_q$.

1. PRELIMINARIES

Throughout this paper, let $q = p^d$ be an arbitrary given power of a prime p , $\mathbb{F}_q$ the finite field of q elements and k the algebraic closure $\bar{\mathbb{F}}_q$ of $\mathbb{F}_q$. For any $n \geq 1$, let $\mathbb{F}_{q^n}$ denote the unique extension field of $\mathbb{F}_q$ of degree n contained in $k = \bar{\mathbb{F}}_q$.

1.1. $\mathbb{F}_q$ -Rational structure on affine varieties.

Definition 1.2. Let (X, A) be an affine variety over $k = \bar{\mathbb{F}}_q$, and $F : X \rightarrow X$ be a morphism. We say that (X, A) is defined over $\mathbb{F}_q$, if the following conditions hold for the algebra homomorphism $F^* : A \rightarrow A$:

- (a) F^* is injective and $F^*(A) = A^q$.
- (b) For each $f \in A$, there exists some $m \geq 1$ such that $(F^*)^m(f) = f^{q^m}$.

In this case, we also say that (X, A) admits an $\mathbb{F}_q$ -rational structure, and that F is the corresponding Frobenius map.

Furthermore, we set $X^F := \{x \in X \mid F(x) = x\}$ for the fixed-point set of F . We also write this set as $X(\mathbb{F}_q)$ and call it the set of $\mathbb{F}_q$ -rational points in X .

In Definition , since $F^*(A) = A^q$ and F^* is injective, we obtain a well-defined map $\sigma : A \rightarrow A$ such that $\sigma(f) = (F^*)^{-1}(f^q)$ for all $f \in A$. It is just an arithmetic Frobenius map satisfying the following:

- (a) The map $\sigma : A \rightarrow A$ is a bijective $\mathbb{F}_q$ -algebra homomorphism such that $\sigma(af) = a^q\sigma(f)$, $a \in k, f \in A$.
- (b) For each $f \in A$, there exists some $m \geq 1$ such that $\sigma^m(f) = f$.

Proposition 1.3. ([15, 4.1.3]) *Let (X, A) be an affine variety defined over $\mathbb{F}_q$ with corresponding Frobenius map F . Set $A_0 := \{f \in A \mid \sigma(f) = f\} = \{f \in A \mid F^*(f) = f^q\}$. Then A_0 is a finitely generated $\mathbb{F}_q$ -subalgebra of A such that the natural map $A_0 \otimes_{\mathbb{F}_q} k \rightarrow A$ is an algebra isomorphism. Furthermore, the fixed-point set X^F is finite, and F is a bijective map.*

1.4. Stable variety under a Frobenius map.

2000 *Mathematics Subject Classification.* 17B20, 17B35, 20G15.

The work was done while the first and second authors were visiting the University of Virginia. They would like to thank the University for its hospitality and financial support. The work is also partially supported by the NSF of China (Grant: 10271047) and CSC.

Suppose (X, A) is an affine variety over k which is defined over $\mathbb{F}_q$, and admits corresponding Frobenius map F . Let V be a subvariety of X , not necessary closed. Call V being F -stable if $F(V) \subset V$.

If V is a closed subset of X , then we have the following property when it's F -stable.

Proposition 1.5. ([15, 4.1.5], [9, 3.3]) *Let (X, A) be an affine variety over k defined over $\mathbb{F}_q$ with corresponding Frobenius map F . Retain the above notations and especially keep $A_0 := \{f \in A \mid \sigma(f) = f\} = \{f \in A \mid F^*(f) = f^q\}$. Then for any closed subset $V \subset X$, the following conditions are equivalent.*

- (a) V is F -stable.
- (b) $F(V) = V$.
- (c) The vanishing ideal $\mathbf{I}_A(V)$ of V is generated by elements in A_0 .
- (d) The ideal $\mathbf{I}_A(V)$ has the form $I_0 \otimes k$ where I_0 is an ideal of A_0 .
- (e) $V = \mathbf{V}_X(S)$ for some subset $S \subset A_0$, where $\mathbf{V}_X(S)$ is the algebraic set defined by S .

Thus, if a closed subvariety V is F -stable, then V is defined over $\mathbb{F}_q$, thereby V admits a rational $\mathbb{F}_q$ -structure.

Example 1.6. Consider the standard Frobenius map $F : k^n \rightarrow k^n$. Let $A = k[x_1, \dots, x_n]$. Let $V \subset k^n$ be an algebraic set. Then A_0 defined in Proposition 1.3 is given by $A_0 = \mathbb{F}_q[x_1, \dots, x_n]$. If V is F -stable, then set $A_0[V] := \mathbb{F}_q[x_1, \dots, x_n]/I_0$, where $I_0 = \mathbf{I}(V) \cap \mathbb{F}_q[x_1, \dots, x_n]$. Then $A_0[V]$ is precisely the subalgebra defined in Proposition 1.3 for $(V, A[V])$.

The following lemma is basic to us.

Theorem 1.7. [24] *Let G be an affine algebraic group, and $F : G \rightarrow G$ a Frobenius map. Suppose X be a G -variety with G -transitive action and with a Frobenius map $F' : X \rightarrow X$, which is compatible with F . This is to say, $F'(g.x) = F(g).F'(x)$ for all $g \in G$ and $x \in X$. Then X admits a fixed point under F' .*

Example 1.8. Let G be a classical group which is a closed subgroup of $GL(m, k)$ and $\mathfrak{g} = \text{Lie}G$. Let F be a standard Frobenius map $F : \mathfrak{gl}(m, k) \rightarrow \mathfrak{gl}(m, k)$ which raises all matrix entries to the q th power. Then the restriction of F are a Frobenius map in G and in $\mathfrak{g} = \text{Lie}(G)$ respectively. In particular, both F are compatible as the above lemma says when G acts on a G -variety in $\mathfrak{g}$ under adjoint action. Recall that any nilpotent orbit $\mathcal{O}$ in $\mathfrak{g}$ has a representative J corresponding to a partition $\mathbf{d} = (d_1 \geq d_2 \geq \dots \geq d_r) \vdash m$ such that J admits ordered Jordan blocks $(J_{d_1}, \dots, J_{d_r})$. Then $J \in \mathfrak{g}^F$. Denote $\mathcal{O}$ by $\mathcal{O}_{\mathbf{d}}$. According to the above lemma, $\mathcal{O}_{\mathbf{d}}$ is F -stable. Notice that

$$(1.8.1) \quad \bar{\mathcal{O}}_{\mathbf{d}} = \cup_{\mathbf{a} \in \{\text{all partitions corresponding to nilpotent orbits in } \text{Lie}(G) \leq \mathbf{d}\}} \mathcal{O}_{\mathbf{a}}.$$

Hence $\bar{\mathcal{O}}$ is also F -stable. Hence, $\bar{\mathcal{O}}$ admits a rational $\mathbb{F}_q$ -structure.

2. FROBENIUS-LIE MORPHISMS AND REPRESENTATIONS OF FINITE RESTRICTED LIE ALGEBRAS

We first recall the definition of Frobenius mappings and $\mathbb{F}_q$ -structures on vector spaces (comparing with the arithmetic Frobenius map in 1.1).

Definition 2.1. (1) Let V and W be two vector space over k . A mapping f from V to W is called a q -semi-linear if it satisfies the following conditions:

- (a) f is an $\mathbb{F}_q$ -linear;
- (b) $f(av) = a^q f(v)$ for all $v \in V$ and $a \in k$.

Furthermore, f is called a q -semi-linear isomorphism if it's bijective. If furthermore $V = W$, then f is called a q -semi-linear automorphism.

(2) A q semi-linear automorphism F on V satisfying that for any $v \in V$, $F^n(v) = v$ for some $n > 0$ is called a q -Frobenius mapping (a Frobenius mapping for short if the q is clear in the context). The smallest r satisfying $F^r(v) = v$ is called a period of v (under F), denoted by $p_F(v)$.

Remark 2.2. (1) It's the same thing both to give a Frobenius mapping F on V and to give an $\mathbb{F}_q$ -structure V_0 on V which means by definition that the natural homomorphism $V_0 \otimes_{\mathbb{F}_q} k \rightarrow V$ is an isomorphism (also, cf. [9, 3.5]). They are related via $V_0 = V^F := \{v \in V \mid F(v) = v\}$. We shall always identify V with $V_0 \otimes k$ in the sequel.

(2) Regard V as a k -vector space variety. Then the coordinate algebra of this variety is the symmetric algebra on V^* : $\mathcal{S}(V^*)$. It's easily seen that F is a morphism of variety.

2.3. Intertwining operator $F_{(V,W)}$.

Let $\mathcal{V}_{k,\mathbb{F}_q}$ be the category whose objects are vector spaces over k with *fixed* $\mathbb{F}_q$ -structures and whose morphisms are linear maps $f : V \rightarrow W$ defined over $\mathbb{F}_q$, namely, $f(V_0) \subseteq W_0$. Clearly, if F_V and F_W are the Frobenius maps on V and W defining V_0 and W_0 , respectively, then f is defined over $\mathbb{F}_q$ if and only if $F_W \circ f = f \circ F_V$. In particular, if F and F' are two Frobenius maps on V , then any $\mathbb{F}_q$ -linear isomorphism from V^F to $V^{F'}$ induces a k -linear isomorphism θ on V such that $\theta \circ F = F' \circ \theta$. Thus, up to isomorphism in $\mathcal{V}_{k,\mathbb{F}_q}$, the $\mathbb{F}_q$ -structure on V is unique. Note that $\mathcal{V}_{k,\mathbb{F}_q}$ is an abelian category.

We shall represent the fixed $\mathbb{F}_q$ -structure on a k -space V by a Frobenius map F_V . Thus, the objects V in $\mathcal{V}_{k,\mathbb{F}_q}$ are, sometimes, written as (V, F_V) , and

$$\mathrm{Hom}_{\mathcal{V}_{k,\mathbb{F}_q}}(V, W) = \{f \in \mathrm{Hom}_k(V, W) \mid F_W \circ f = f \circ F_V\}.$$

For $(V, F_V), (W, F_W) \in \mathrm{Ob}(\mathcal{V}_{k,\mathbb{F}_q})$, there is an induced $\mathbb{F}_q$ -linear map

$$F_{(V,W)} : \mathrm{Hom}_k(V, W) \rightarrow \mathrm{Hom}_k(V, W); f \mapsto F_{(V,W)}(f) = F_W \circ f \circ F_V^{-1}.$$

Note that, when V is *infinite dimensional*, $F_{(V,W)}$ is not necessarily a Frobenius map on $\mathrm{Hom}_k(V, W)$. However, we have the following ([8]).

Lemma 2.4. *Maintain the notation introduced above and let $F = F_{(V,W)}$. We have an $\mathbb{F}_q$ -space isomorphism*

$$\mathrm{Hom}_{\mathbb{F}_q}(V^{F_V}, W^{F_W}) \cong \mathrm{Hom}_{\mathcal{V}_{k,\mathbb{F}_q}}(V, W) = \mathrm{Hom}_k(V, W)^F.$$

In particular, if V is finite dimensional, then F is a Frobenius map on $\mathrm{Hom}_k(V, W)$.

2.5. Frobenius-Lie morphisms.

Let $\mathfrak{g}$ be a finite-dimensional Lie algebra over k . A mapping $F : \mathfrak{g} \rightarrow \mathfrak{g}$ is called a *Frobenius-Lie morphism* on $\mathfrak{g}$ if it is a Frobenius mapping on the k -space $\mathfrak{g}$, and if it preserves the Lie product: $F[x, y] = [F(x), F(y)]$ for all $x, y \in \mathfrak{g}$. Clearly, a Frobenius-Lie morphism on $\mathfrak{g}$ is also a Lie algebra $\mathbb{F}_q$ -automorphism. Also, if $\mathfrak{g} = A^-$ is the Lie algebra obtained from an associative algebra A with the Lie product $[x, y] = xy - yx$ for all $x, y \in A$, then any Frobenius morphism on A induces a Frobenius-Lie morphism on A^- .

The set of all Frobenius-Lie morphisms on $\mathfrak{g}$ is denoted by $\mathfrak{F}_q(\mathfrak{g})$. For any $\sigma \in \mathrm{Aut}(\mathfrak{g})$, it's easy to see by direct verification that $\sigma_* F := \sigma^{-1} F \sigma$ is still a Frobenius-Lie morphism on $\mathfrak{g}$. Two Frobenius-Lie morphisms F_1 and F_2 on $\mathfrak{g}$ are called being equivalent, denoted by $F_1 \sim F_2$, if $\mathfrak{g}^{F_1} \cong \mathfrak{g}^{F_2}$ as Lie $\mathbb{F}_q$ -algebras. The following Lemma is clear ([10]).

Lemma 2.6. *Let F_1 and F_2 be two Frobenius Lie morphisms on $\mathfrak{g}$.*

- (a) $F_1 = F_2$ if and only if $\mathfrak{g}^{F_1} = \mathfrak{g}^{F_2}$;
- (b) $F_1 \sim F_2$ if and only if $F_1 = \sigma_*(F_2)$ for some $\sigma \in \mathrm{Aut}(\mathfrak{g})$.

2.7. $F_{\mathfrak{g}}$ -stable modules.

Let V be a finite dimensional $\mathfrak{g}$ -module and let $\rho = \rho_V : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ be the corresponding representation. We say that V is $F_{\mathfrak{g}}$ -stable (or simply F -stable) if there is an $\mathbb{F}_q$ -structure V_0 of V such that ρ induces a representation $\rho_0 : \mathfrak{g}^F \rightarrow \mathfrak{gl}_{\mathbb{F}_q}(V_0)$ of $\mathfrak{g}^F$. By the definition of an $\mathbb{F}_q$ -structure, V is F -stable if and only if there is a Frobenius map $F_V : V \rightarrow V$ such that

$$(2.7.1) \quad F_V(xv) = F_{\mathfrak{g}}(x)F_V(v), \quad \text{for all } x \in \mathfrak{g}, v \in V.$$

In terms of the corresponding representation ρ , the F -stability of V simply means that there is a Frobenius map F_V on V such that $\rho \circ F_{\mathfrak{g}} = F_{(V,V)} \circ \rho$. Thus, if V is an F -stable $\mathfrak{g}$ -module with respect to F_V , then $V = V^F \otimes_{\mathbb{F}_q} k$, $F_V(v \otimes a) = v \otimes a^q$ for all $v \in V^F$, $a \in k$, and V^F is an $\mathfrak{g}^F$ -module. Here, again for notational simplicity, we write V^F for V^{F_V} . What's more, we sometimes use $\mathfrak{g}^F$ and V^F in the meantime, but obviously with different meanings of both F 's.

Up to isomorphism, it doesn't matter which Frobenius maps (or $\mathbb{F}_q$ -structures) on V we shall work with when considering F -stable modules. This is because we have the following lemma which directly follows from Noether-Deuring Theorem (c.f. [7, p.139]).

Lemma 2.8. *Let (V_1, F_1) and (V_2, F_2) be two F -stable $\mathfrak{g}$ -modules. Then $V_1^{F_1} \cong V_2^{F_2}$ as $\mathfrak{g}^F$ -modules if and only if $V_1 \cong V_2$ as $\mathfrak{g}$ -modules. In particular, if $V_1 = V_2 = V$, then $V^{F_1} \cong V^{F_2}$ as $\mathfrak{g}^F$ -modules.*

In the following, we sometimes use the notation (V, F_V) for an F -stable module V . Let $\mathbf{mod}^F\text{-}\mathfrak{g}$ denote the category whose objects are F -stable modules $V = (V, F_V)$. The morphisms from (V, F_V) to (W, F_W) are given by homomorphisms $\theta \in \text{Hom}_{\mathfrak{g}}(V, W)$ such that $\theta \circ F_V = F_W \circ \theta$, that is, $\mathfrak{g}$ -module homomorphisms compatible with their $\mathbb{F}_q$ -structures. It is easy to see that $\mathbf{mod}^F\text{-}\mathfrak{g}$ is an abelian $\mathbb{F}_q$ -category. The following result allows us to embed a module category over a finite field into a category over the algebraic closure of the finite field.

Theorem 2.9. ([10]) *Let $\mathfrak{g}$ is a finite-dimensional Lie algebra over k . The abelian category $\mathbf{mod}^F\text{-}\mathfrak{g}$ is equivalent to the category $\mathbf{mod}\text{-}\mathfrak{g}^F$ of finite dimensional $\mathfrak{g}^F$ -modules. In particular, there is a one-to-one correspondence between isoclasses of irreducible (indecomposable, resp.) $\mathfrak{g}^F$ -modules and isoclasses of irreducible (indecomposable, resp.) F -stable $\mathfrak{g}$ -modules.*

Remark 2.10. In general, a $\mathfrak{g}$ -module may not be F -stable. But each finite-dimensional module V must be $F^{n(V)}$ -stable for a certain $n(V) \in \mathbb{N}$. Thus, for a given finite-dimensional module V there is a so-called F -orbit of V , arising from V under some twist action of $\mathfrak{g}$. The sum of twisted modules running through the orbit is F -stable. Furthermore, there is a one-to-one correspondence between the isoclasses of indecomposable (resp. irreducible) $\mathfrak{g}^F$ -modules and the F -orbits of the isoclasses of indecomposable (resp. irreducible) $\mathfrak{g}$ -modules ([10]).

2.11. Finite restricted Lie algebras.

From this section on, we will focus our discussion on restricted Lie algebras. Always suppose the Lie algebra $\mathfrak{g}$ is endowed with a p -mapping $[p]$. Suppose F is still a Frobenius-Lie morphism. Call F a restricted Frobenius morphism if it satisfies an additional condition $F(x^{[p]}) = F(x)^{[p]}$.

Lemma 2.12. *Suppose F is a Frobenius-Lie morphism on a restricted Lie algebra $(\mathfrak{g}, [p])$. Then*

- (a) *F is a restricted morphism if and only if $\mathfrak{g}^F$ is a restricted subalgebra of $\mathfrak{g}$ over $\mathbb{F}_q$.*
- (b) *If F is a restricted morphism, then $(q^n\text{-})$ Frobenius-Lie morphism F^n is also restricted.*

3. PROJECTIVENESS AND EXTENSIONS FOR RESTRICTED F -STABLE MODULES

We always suppose in this section that $\mathfrak{g}$ is a restricted Lie algebra with a Frobenius-Lie morphism $F_{\mathfrak{g}}$.

3.1. Projectiveness

Lemma 3.2. *Let $\mathfrak{g}$ be a finite dimensional restricted Lie algebra over k , with a restricted Frobenius-Lie morphism $F_{\mathfrak{g}}$, and $\mathfrak{g}_0 = \mathfrak{g}^{F_{\mathfrak{g}}}$. Suppose (P, F) is an $F_{\mathfrak{g}}$ -stable $\mathfrak{g}$ -module (or simply: F -stable $\mathfrak{g}$ -module), and $P_0 = P^F$. Then P is a projective $\mathfrak{g}$ -module if and only if P_0 is a projective $\mathfrak{g}_0$ -module.*

Proof. (1) It follows from the exactness of the field extension functor that the projectiveness of P_0 implies the projectiveness of P .

(2) Now assume that P is a projective $\mathfrak{g}$ -module. In order to verify the projectiveness of P_0 , it's sufficient to prove that any short exact sequence of $\mathfrak{g}_0$ -modules

$$(3.2.1) \quad 0 \rightarrow M_0 \rightarrow N_0 \rightarrow P_0 \rightarrow 0$$

is split. Set $M = M_0 \otimes_{\mathbb{F}_q} k$, and $N = N_0 \otimes_{\mathbb{F}_q} k$. Then we have a short exact sequence of $\mathfrak{g}$ -modules

$$(3.2.2) \quad 0 \rightarrow M \rightarrow N \rightarrow P \rightarrow 0.$$

The projectiveness of P implies that the short exact sequence (3.2.2) is split. Hence we have $N \cong P \oplus M$, as $\mathfrak{g}$ -modules. Owing to Noether-Deuring Theorem (cf. [7, p.139]), $N_0 \cong M_0 \oplus P_0$. Thus the exact sequence (3.2.1) is split. Hence P_0 is really projective. We are done. $\square$

Proposition 3.3. *Suppose $\mathfrak{g}$ is a restricted Lie algebra with a restricted Frobenius-Lie morphism $F_{\mathfrak{g}}$. If all restricted simple modules of $\mathfrak{g}$ are F -stable, then all restricted projective $\mathfrak{g}$ -modules are F -stable. Especially, for $\mathfrak{g}$ a classical Lie algebra in sense of [23], all restricted projective modules of $\mathfrak{g}$ are F -stable under any given Frobenius-Lie morphism $F_{\mathfrak{g}}$.*

Proof. Consider restricted $u(\mathfrak{g})$ -module category $\mathbf{mod}\text{-}u(\mathfrak{g})$ and $u(\mathfrak{g}_0)$ -module category $\mathbf{mod}\text{-}u(\mathfrak{g}_0)$. Suppose (S, F_S) is a simple F -stable $u(\mathfrak{g})$ -module. Set $S_0 = S^{F_S}$, which is a simple $u(\mathfrak{g}_0)$ -module (cf. Theorem 2.9). For the projective cover $P(S_0)$ in $\mathbf{mod}\text{-}u(\mathfrak{g}_0)$, set $P(S_0)_k := P(S_0) \otimes_{\mathbb{F}_q} k$. Then $P(S_0)_k$ is a projective cover of S , thereby it is an indecomposable projective $\mathfrak{g}$ -module (cf. Theorem 2.9). Recall that indecomposable projective $u(\mathfrak{g})$ -modules (resp. $u(\mathfrak{g}_0)$ -modules) are in one-to-one correspondence to irreducible $u(\mathfrak{g})$ -modules (resp. $u(\mathfrak{g}_0)$ -modules) (cf. [7]). By Lemma 3.2, all indecomposable projective $u(\mathfrak{g})$ -modules are F -stable provided that all restricted simple modules of $\mathfrak{g}$ are F -stable. Thus, all projective $u(\mathfrak{g})$ -modules are F -stable as long as all restricted simple modules of $\mathfrak{g}$ are F -stable.

As to the second statement concerning the classical Lie algebra $\mathfrak{g}$, we only need to prove that all simple restricted $\mathfrak{g}$ -modules are F -stable. Recall from [10, 5.2] that any Frobenius-Lie morphism $F_{\mathfrak{g}}$ on $\mathfrak{g}$ is of the form $F_{\mathfrak{g}} = \gamma \cdot g^{-1} F_0 g$ for $\gamma \in \Gamma$ a graph automorphism of the Dynkin diagram, and for $g \in G$ which coincides with the Chevalley group $G(\mathfrak{h})$ constructed relative to $\mathfrak{h}$ (cf. [23, IV.6]). Here F_0 is a split Frobenius-Lie morphism, related to $\mathfrak{h}$, i.e. the fixed-point space of F_0 is just the split $\mathbb{F}_q$ -form of $\mathfrak{g}$. Hence $F_{\mathfrak{g}}$ is restricted, thereby $\mathfrak{g}_0 := \mathfrak{g}^{F_{\mathfrak{g}}}$ is a restricted Lie subalgebra of $\mathfrak{g}$. Note that there is an $h \in G$ such that $\gamma \cdot F_0 = h F_{\mathfrak{g}} h^{-1}$ ([10, Theorem 5.4]). We may assume $F_{\mathfrak{g}} = \gamma \cdot F_0$ in the following argument for simplicity, without a loss of universality. Recall that $\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}$, where $\mathfrak{h}$ is a given Cartan subalgebra, $\mathfrak{n}$ (resp. $\mathfrak{n}^-$) is the sum of positive root spaces (resp. the sum for negative root spaces), associated with $\mathfrak{h}$. Set $\mathfrak{b} := \mathfrak{n} \oplus \mathfrak{h}$, a Borel subalgebra of $\mathfrak{g}$. Hence $\mathfrak{n}$, $\mathfrak{h}$, $\mathfrak{n}^-$ and $\mathfrak{b}$ are all $F_{\mathfrak{g}}$ -stable. Correspondingly, we have the triangular decomposition $\mathfrak{g}_0 = \mathfrak{n}_0^- \oplus \mathfrak{h}_0 \oplus \mathfrak{n}_0$, and $\mathfrak{b}_0 = \mathfrak{h}_0 \oplus \mathfrak{n}_0$ with $\mathfrak{h}_0 = \mathfrak{h}^{F_{\mathfrak{g}}}$, $\mathfrak{n}_0^- = (\mathfrak{n}^-)^{F_{\mathfrak{g}}}$, $\mathfrak{n}_0 = \mathfrak{n}^{F_{\mathfrak{g}}}$ and $\mathfrak{b}_0 = \mathfrak{b}^{F_{\mathfrak{g}}}$. Notice that irreducible restricted $\mathfrak{g}$ -modules are parameterized by $\Lambda := \{\lambda \in \mathfrak{h}^* \mid \lambda(h)^p = \lambda(h) \text{ for } h \in \mathfrak{h}_0\}$ as follows: each simple module $L(\lambda)$ for $\lambda \in \Lambda$ is the unique quotient of the baby Verma module $Z(\lambda)$ by the unique maximal submodule M in $Z(\lambda)$. Here $Z(\lambda) = u(\mathfrak{g}) \otimes_{u(\mathfrak{b})} k_{\lambda}$, for the one dimensional module k_{λ} with $\mathfrak{h}$ -action by multiplicity λ and trivial $\mathfrak{n}$ -action. And M is the sum of all proper submodules which can be described by not containing the $u(\mathfrak{g})$ -generator $1 \otimes 1$ in $Z(\lambda)$. Also, all simple restricted $\mathfrak{g}_0$ -modules can be parameterized by a counterpart $L_0(\lambda)$, λ in $\Lambda_0 = \{\lambda \in \mathfrak{h}_0^* \mid \lambda(h) \in \mathbb{F}_p \text{ for } h \in \mathfrak{h}_0\}$. Obviously, Λ_0 can be identified with Λ . Furthermore,

$L_0(\lambda) = Z_0(\lambda)/M_0$ for $Z_0(\lambda)_0 = u(\mathfrak{g}_0) \otimes_{u(\mathfrak{b}_0)} (\mathbb{F}_q)_\lambda$ with the same meaning as for $Z(\lambda)$, M_0 is the unique maximal submodule of $Z(\lambda)_0$. Those $L(\lambda)$ just have $\mathbb{F}_q$ -structures $L(\lambda)_0$. Hence all of them are F -stable. $\square$

3.4. Extensions.

Let us first recall of the following fact.

Lemma 3.5. *Let $\mathfrak{g}$ be a restricted Lie algebra over k , (V, F_V) and (W, F_W) two given F -stable restricted $\mathfrak{g}$ -modules. Set $V_0 = V^{F_V}$ and $W_0 = W^{F_W}$. Then $\text{Hom}_{\mathfrak{g}}(V, W)^{F_{(V,W)}} \cong \text{Hom}_{\mathfrak{g}_0}(V_0, W_0)$, where the Frobenius mapping $F_{(V,W)}$ sends ϕ to $F_V \circ \phi \circ F_W^{-1}$.*

Proof. It can be verified by straightforward computation that $F_{(V,W)}$ preserves $\mathfrak{g}$ -module homomorphisms. Then it follows from Lemma 2.4. $\square$

In the following, we will see that the above isomorphism can be extended to the higher extensions between V and W . Set $\mathbf{P}(V_0)$ be a projective resolution of V_0 in $\mathbf{mod}\text{-}u(\mathfrak{g}_0)$, where $\mathbf{P}(V_0)$ is as follows:

$$\mathbf{P}(V_0) : \cdots \longrightarrow P_0^{(n)} \xrightarrow{\delta_n} P_0^{(n-1)} \xrightarrow{\delta_{n-1}} \cdots \xrightarrow{\delta_1} P_0^{(0)} \xrightarrow{\delta_0} V_0.$$

Set the contravariant functor $\Gamma_0 = \text{Hom}_{u(\mathfrak{g}_0)}(-, W_0)$. We have a complex $(\Gamma_0(\mathbf{P}(V_0)), \delta^*)$, where $\delta_n^* : \Gamma_0(P_0^{n-1}) \rightarrow \Gamma_0(P_0^n)$ with $\delta_n^*(\phi) = \phi \circ \delta_n$. And we have $\text{Ext}_0^n(V_0, W_0) = R^n \Gamma_0(\mathbf{P}(V_0))$. Now arising from the field extension $\mathbb{F}_q \hookrightarrow k$, there is a projective resolution of the $u(\mathfrak{g})$ -module V

$$\mathbf{P}(V) : \cdots \longrightarrow P^{(n)} \xrightarrow{\delta_n \otimes 1} P^{(n-1)} \xrightarrow{\delta_{n-1} \otimes 1} \cdots \xrightarrow{\delta_1 \otimes 1} P^{(0)} \xrightarrow{\delta_0 \otimes 1} V$$

where $P^{(n)} = P_0^{(n)} \otimes_{\mathbb{F}_q} k$ with natural Frobenius mappings denoted by F_n . The contravariant functor $\Gamma = \text{Hom}_{u(\mathfrak{g})}(-, W)$ gives rise to the cohomology of the complex $\Gamma(\mathbf{P}(V))$: $\text{Ext}^*(V, W)$. Recall that

$$\text{Ext}^n(V, W) = \{\bar{\phi} \in \text{Ker}(\delta_{n+1} \otimes 1)^* / \text{Im}(\delta_n \otimes 1)^* \mid \phi \in \text{Ker}(\delta_{n+1} \otimes 1)^*\}.$$

Notice that $(\delta_n \otimes 1) \circ F_n = F_{n-1} \circ (\delta_n \otimes 1)$. It's easily verified by direct computation that both $F_{(P_n, W)}|_{\text{Im}(\delta_n \otimes 1)^*}$ and $F_{(P_n, W)}|_{\text{Ker}(\delta_{n+1} \otimes 1)^*}$ are Frobenius mappings in the respective vector spaces. Define $\mathfrak{F}_n : \text{Ext}^n(V, W) \rightarrow \text{Ext}^n(V, W)$ by $\mathfrak{F}_n(\bar{\phi}) = \overline{F_{(P_n, W)}(\phi)}$. Then $\mathfrak{F}_n$ is really a Frobenius mapping of $\text{Ext}^n(V, W)$, whose fixed point space is just $\text{Ext}_0^n(V_0, W_0)$. Hence we have

Proposition 3.6. *Suppose $\mathfrak{g}$ is a restricted Lie algebra with a Frobenius-Lie morphism $F_{\mathfrak{g}}$. Suppose (V, F_V) and (W, F_W) are two finite-dimensional restricted F -stable $\mathfrak{g}$ -modules with $V_0 = V^{F_V}$ and $W_0 = W^{F_W}$. Denote by $\text{Ext}^n(V, W)$ (resp. $\text{Ext}_0^n(V_0, W_0)$) the n -th extension between restricted $\mathfrak{g}$ -modules V and W over $u(\mathfrak{g})$ (resp. the n -th extension for $u(\mathfrak{g}_0)$ -modules V_0 and W_0). Then $\text{Ext}^n(V, W)^{\mathfrak{F}_n} \cong \text{Ext}_0^n(V_0, W_0)$ for $n = 0, 1, 2, \dots$.*

Next we will give a corollary which will be useful in the next section. Before this, we need the following obvious fact.

Lemma 3.7. *Let (A, F_A) be a finite-dimensional associative algebra over k with a Frobenius morphism. (V, F_M) is a finite-dimensional F_A -stable A -module (which has the same meaning as $F_{\mathfrak{g}}$ -stable modules, cf. [8]). Set $A_0 = A^{F_A}$ and $M_0 = M^{F_M}$. Then $\text{Ann}_A(M) = \text{Ann}_{A_0}(M_0) \otimes_{\mathbb{F}_q} k$.*

Corollary 3.8. *Maintain the same notations as in Proposition 3.6. Then*

$$\text{Ann}_{H^{ev}(u(\mathfrak{g}), k)}(\text{Ext}^{ev}(M, M)) = \text{Ann}_{H^{ev}(u(\mathfrak{g}_0), \mathbb{F}_q)}(\text{Ext}_0^{ev}(M_0, M_0)) \otimes_{\mathbb{F}_q} k.$$

4. $\mathbb{F}_q$ -RATIONAL STRUCTURE ON SUPPORT VARIETIES

4.1. We recall some facts about support varieties as follows (cf. [11], [12] and [13]).

For any restricted Lie algebra $\mathfrak{g}$ one has the cohomology variety $|\mathfrak{g}|$ of $\mathfrak{g}$ which is the affine variety associated to $H^{\text{ev}}(u(\mathfrak{g}), k)^{(-1)}$. Here and further, the $H^{\text{ev}}(u(\mathfrak{g}), k)^{(-1)}$ with superscript (-1) means the twisted ring. Let V be a finite-dimensional $\mathfrak{g}$ -module, the support variety $|\mathfrak{g}|_V$ is the support in $|\mathfrak{g}|$ of the $H^{\text{ev}}(u(\mathfrak{g}), k)^{(-1)}$ -module $H^\bullet(u(\mathfrak{g}), V \otimes V^*)^{(-1)}$, $|\mathfrak{g}|_V := \text{Supp}(H^\bullet(u(\mathfrak{g}), V \otimes V^*)^{(-1)})$ (cf. [12], [13]). Denote by $\Phi : |\mathfrak{g}| \rightarrow \mathfrak{g}$ the morphism of varieties associated with the natural map $\Phi^* : S^\bullet(\mathfrak{g}^*) \rightarrow H^{\text{ev}}(u(\mathfrak{g}), k)^{(-1)}$, where $S^\bullet(\mathfrak{g}^*)$ is the symmetric algebra on the dual of $\mathfrak{g}$. We may identify $\Phi(|\mathfrak{g}|_V)$ with the support of the graded $S^\bullet(\mathfrak{g}^*)$ -module $H^\bullet(u(\mathfrak{g}), V \otimes V^*)^{(-1)}$. So defined, $\Phi(|\mathfrak{g}|_V)$ is a conical closed subvariety of the restricted nilpotent cone $\mathcal{N}_1 := \{x \in \mathfrak{g} \mid x^{[p]} = 0\}$: $\Phi(|\mathfrak{g}|_V) = \{x \in \mathcal{N}_1 \mid V \text{ is not projective as a restricted } \langle x \rangle\text{-module}\} \cup \{0\}$ (cf. [18] and [12]). We will often omit the superscript (-1) in the following argument.

Lemma 4.2. *Suppose $\mathfrak{g}$ is a restricted Lie algebra over k , defined over $\mathbb{F}_q$ with respect to Frobenius-Lie morphism $F_{\mathfrak{g}}$ on $\mathfrak{g}$. Then $\Phi(|\mathfrak{g}|)$ is defined over $\mathbb{F}_q$ such that the associated $\mathbb{F}_q$ -rational structure is given by the restriction of $F_{\mathfrak{g}}$ in $\Phi(|\mathfrak{g}|)$.*

Proof. Recall that $\Phi(|\mathfrak{g}|) = \mathcal{N}_1(\mathfrak{g})$. Regard $\Phi(|\mathfrak{g}|)$ as a closed variety of k -vector space variety $\mathfrak{g}$. According to the argument in the proof of Proposition 3.3, $F_{\mathfrak{g}}$ is restricted. Hence $F_{\mathfrak{g}}(\Phi(|\mathfrak{g}|)) \subset \Phi(|\mathfrak{g}|)$. By Remark 2.2 and Proposition 1.5, $F_{\mathfrak{g}}$ is a morphism of variety, and $\Phi(|\mathfrak{g}|)$ is defined over $\mathbb{F}_q$ with the restriction of $F_{\mathfrak{g}}$, as the corresponding Frobenius map. $\square$

Suppose V_0 is a restricted $\mathfrak{g}_0$ -module. Recall of the construction of a minimal projective resolution $\cdots \rightarrow P_0^{(n)} \rightarrow \cdots \rightarrow P_0^{(0)} \rightarrow V_0$ of V_0 . Set $P_0^{(0)} = P(V_0)$, the projective cover of V_0 , and define the syzygy module $\Omega(V_0)$ as the kernel of the given map $P(V_0) \rightarrow V_0$. Inductively, set $P_0^{(n)} = P(\Omega^n(V_0))$ and $\Omega^n(V_0) = \Omega(\Omega^{n-1}(V_0))$. When $V_0 = \mathbb{F}_q$, then $H^n(u(\mathfrak{g}_0), \mathbb{F}_q) \cong \text{Hom}_{u(\mathfrak{g}_0)}(\Omega^n(\mathbb{F}_q), \mathbb{F}_q)$. Suppose $\zeta \in H^{2n}(u(\mathfrak{g}_0), \mathbb{F}_q)$. Such a ζ can be represented by a map $\zeta : \Omega^{2n}(\mathbb{F}_q) \rightarrow \mathbb{F}_q$ whose kernel is the $u(\mathfrak{g}_0)$ -module $\mathbb{L}_{\zeta}$. In the meantime, we have for $V = V_0 \otimes_{\mathbb{F}_q} k$, $P^{(n)} = P(\Omega^n(V))$ and $\Omega^n(V) = \Omega(\Omega^{n-1}(V))$. By Proposition 3.3 and the exactness of the field extension functor from the $u(\mathfrak{g}_0)$ -module category to the $u(\mathfrak{g})$ -module category, $P^{(n)} \cong P_0^{(n)} \otimes_{\mathbb{F}_q} k$, and $\Omega^n(V) \cong \Omega^n(V_0) \otimes_{\mathbb{F}_q} k$. Especially, $\Omega^{2n}(k) \cong \Omega^{2n}(\mathbb{F}_q) \otimes_{\mathbb{F}_q} k$.

Lemma 4.3. *Maintain the notations as above. Set $L_{\zeta} = \mathbb{L}_{\zeta} \otimes_{\mathbb{F}_q} k$, which is a restricted $\mathfrak{g}$ -module. Then L_{ζ} is the kernel of $\zeta \otimes 1 : \Omega^{2n}(k) \rightarrow k$.*

Proof. We only need to prove that $\ker(\zeta \otimes 1) = \ker(\zeta) \otimes_{\mathbb{F}_q} k$. Actually, any short exact sequence of $u(\mathfrak{g}_0)$ -modules $S_1 \hookrightarrow S_2 \twoheadrightarrow S_3$ gives rise to the short exact sequence of $u(\mathfrak{g})$ -modules: $T_1 \hookrightarrow T_2 \twoheadrightarrow T_3$ for $T_i = S_i \otimes_{\mathbb{F}_q} k$, $i = 1, 2, 3$. This fact implies that $\ker(\zeta \otimes 1) = \ker(\zeta) \otimes_{\mathbb{F}_q} k$. $\square$

Theorem 4.4. *Maintain the notations as in Lemma 4.2. If V is a restricted $F_{\mathfrak{g}}$ -stable $\mathfrak{g}$ -module, then there is an $\mathbb{F}_q$ -rational structure on $\Phi(|\mathfrak{g}|_V)$. Conversely, if X is a closed conical subvariety of $\Phi(|\mathfrak{g}|)$ which admits an $\mathbb{F}_q$ -rational structure arising from $F_{\mathfrak{g}}$, then there exists a restricted $F_{\mathfrak{g}}$ -stable $\mathfrak{g}$ -module V such that $X = \Phi(|\mathfrak{g}|_V)$.*

Proof. (1) Suppose (V, F_V) is an F -stable restricted $\mathfrak{g}$ -module, $V_0 = V^{F_V}$. We will prove that $\Phi(|\mathfrak{g}|_V)$ is really defined over $\mathbb{F}_q$. Recall $\Phi(|\mathfrak{g}|_V)$ is a closed subvariety of $\Phi(|\mathfrak{g}|)$ which is defined over $\mathbb{F}_q$ (Lemma 4.2). Furthermore, $H^{\text{ev}}(u(\mathfrak{g}), k) \cong H^{\text{ev}}(u(\mathfrak{g}_0), \mathbb{F}_q) \otimes_{\mathbb{F}_q} k$ (Lemma 4.2). By Proposition 1.5, it's sufficient to prove that the defining ideal of $\Phi(|\mathfrak{g}|_V)$ is of the form $I_0 \otimes_{\mathbb{F}_q} k$ where I_0 is an ideal of $H^{\text{ev}}(u(\mathfrak{g}_0), \mathbb{F}_q)$. This follows from Lemma 3.8.

(2) Suppose X is a closed conical subvariety of $\Phi(|\mathfrak{g}|)$ defined over $\mathbb{F}_q$. We will construct a restricted F -stable $\mathfrak{g}$ -module V such that $X = \Phi(|\mathfrak{g}|_V)$. Recall that in [12], any closed conical

subvariety of $\Phi(|\mathfrak{g}|)$ can be realized as a support variety of a restricted $\mathfrak{g}$ -modules, which follows from the Carlson's construction in [4]. What we need to do in our case is to prove that such a construction of modules can be defined over $\mathbb{F}_q$ (cf. [1, Theorem, 2.26.9]).

As X is closed and conical, X is defined by a homogenous ideal I in $H^{\text{ev}}(u(\mathfrak{g}), k)$. Since $H^{\text{ev}}(u(\mathfrak{g}), k)$ is Noetherian (cf. [11]), I is finitely generated by some homogeneous elements $I = \langle \zeta_1, \dots, \zeta_r \rangle$. Furthermore, by Proposition 3.6 $H^{\text{ev}}(u(\mathfrak{g}), k)$ has an $\mathbb{F}_q$ -form $H_0^{\text{ev}}(u(\mathfrak{g}_0), \mathbb{F}_q)$, i.e.

$$H^{\text{ev}}(u(\mathfrak{g}), k) \cong H_0^{\text{ev}}(u(\mathfrak{g}_0), \mathbb{F}_q) \otimes_{\mathbb{F}_q} k,$$

as k -algebras. By assumption X is defined over $\mathbb{F}_q$ which means that I is an ideal generated by elements of $H_0^{\text{ev}}(u(\mathfrak{g}_0), \mathbb{F}_q)$ (Proposition 1.5). Thus those ζ_i can be chosen in $H_0^{\text{ev}}(u(\mathfrak{g}_0), \mathbb{F}_q)$. Suppose $\zeta_i \in H^{2n}(u(\mathfrak{g}_0), \mathbb{F}_q)$. By Lemma 4.3, there is a $u(\mathfrak{g}_0)$ -submodule $\mathbb{L}_{\zeta_i} \in \Omega^{2n}(\mathbb{F}_q)$ such that $\mathbb{L}_{\zeta_i}$ is just the kernel of $\zeta_i : \Omega^n(\mathbb{F}_q) \rightarrow \mathbb{F}_q$ which represents the cohomology class $\zeta_i \in H^{2n}(u(\mathfrak{g}_0), \mathbb{F}_q) \cong \text{Hom}_{u(\mathfrak{g}_0)}(\Omega^{2n}(\mathbb{F}_q), \mathbb{F}_q)$. And $L_{\zeta_i} = \mathbb{L}_{\zeta_i} \otimes_{\mathbb{F}_q} k$ is just the Carlson's module (cf. [4]). Set $V = L_{\zeta_1} \otimes \dots \otimes L_{\zeta_r}$, then by [12, Theorem 2.2] $X = \Phi(|\mathfrak{g}|_V)$. And this V is $F_{\mathfrak{g}}$ -stable. We are done. $\square$

Let G be an affine algebraic group over k , defined over $\mathbb{F}_q$, and $\mathfrak{g} = \text{Lie}(G)$. We call a rational G -module stable if it's $F_{\mathfrak{g}}$ -stable as a $\mathfrak{g}$ -module. By [12, Theorem 1.2], we have a corollary to the above theorem.

Corollary 4.5. *Suppose V is a finite-dimensional rational stable G -module V , then $\Phi(|\mathfrak{g}|_V)$ is a closed conical G -invariant variety defined over $\mathbb{F}_q$.*

The inverse statement of the above is a version of an old Friedlander-Parshall's conjecture ([12]): any closed conical G -invariant subvariety of $\Phi(|\mathfrak{g}|)$ is the support variety of a finite-dimensional rational stable G -module V .

5. F -STABLE NILPOTENT ORBITS

In the whole section, we assume that $\mathfrak{g}$ is a simple classical Lie algebra over k an algebraically closed field of characteristic $p > 3$, which will be additionally assumed to be good for the root system of $\mathfrak{g}$ in the subsection 5.3 (cf. [5, p.14]).

5.1. Frobenius-Lie morphisms and the graph automorphisms of Dynkin diagrams. Recall that a classical simple Lie algebra is classified by Dynkin diagrams labelled by A_n ($n \geq 1$), B_n ($n \geq 2$), C_n ($n \geq 3$), D_n ($n \geq 4$), E_n ($n = 6, 7, 8$), F_4 , G_2 (see [23, II.10.1]). We may view such a $\mathfrak{g}$ as reduction modulo p via a Chevalley basis of the corresponding complex simple Lie algebra $\mathfrak{g}_{\mathbb{C}}$ except for A_n with $p|n$. In the exceptional cases (i.e. A_n with $p|n$), $\mathfrak{g}$ is a quotient of the reduction modulo p via Chevalley basis by the one-dimensional center which is in Cartan subalgebras. There are naturally resulting basis in $\mathfrak{g}$ arising from the Chevalley basis of $\mathfrak{g}_{\mathbb{C}}$, still called Chevalley basis (to see below for more arguments). Let G be the adjoint group of $\mathfrak{g}$. By Chevalley-Dickson's result, G is simple (cf. [25, §4, Theorem 5] or [23, Theorem 3.1]) with $\mathfrak{g} = \text{Lie}(G)$ for non-exceptional cases, and $\mathfrak{g} = [\text{Lie}(G), \text{Lie}(G)]$ or $\mathfrak{g} = [\text{Lie}(G), \text{Lie}(G)]/\text{1-dimensional center of } [\text{Lie}(G), \text{Lie}(G)]$ for the exceptional case ([16, 5.4]).

We have a realization of G in $\text{Aut}(\mathfrak{g})$ as follows. Denote by $G_{\mathbb{C}} \subset GL(n, \mathbb{C})$ the simply connected simple algebraic group over $\mathbb{C}$ with $\text{Lie}(G_{\mathbb{C}}) = \mathfrak{g}_{\mathbb{C}}$, associated with root system Φ and simple root system. Let $\mathfrak{g}_{\mathbb{C}} = \text{Lie}(G_{\mathbb{C}})$ and let $\mathcal{B} = \{X_{\alpha} \mid \alpha \in \Phi\} \cup \{H_i \mid i = 1, \dots, r = \text{rank}(\mathfrak{g}_{\mathbb{C}})\}$. Here the Cartan subalgebra $\mathfrak{h}_{\mathbb{C}} = \sum_i \mathbb{C}H_i$. We may suppose that H consists of diagonal matrices. Set $\mathfrak{h}_{\mathbb{Z}} = \sum_i \mathbb{Z}H_i$ (resp. $\mathfrak{g}_{\mathbb{Z}} = \sum_{\alpha \in \Phi} \mathbb{Z}X_{\alpha} + \mathfrak{h}_{\mathbb{Z}}$) and $\mathfrak{h}_k = \mathfrak{h}_{\mathbb{Z}} \otimes_{\mathbb{Z}} k$ (resp. $\mathfrak{g}_k = \mathfrak{g}_{\mathbb{Z}} \otimes_{\mathbb{Z}} k$). Note that $\mathfrak{g}_k$ has a Chevalley basis $\mathcal{B}_k = \{e_{\alpha} := X_{\alpha} \otimes 1 \mid \alpha \in \Phi\} \cup \{h_{\alpha_i} := H_{\alpha_i} \otimes 1 \mid i = 1, \dots, r\}$, where $\Pi = \{\alpha_i \mid i = 1, 2, \dots, r\}$ is the simple root set of Φ .

Let $\bar{G}$ be the simply connected simple algebraic group over k , associated with the irreducible root system Φ . Then $\text{Lie}(\bar{G}) = \mathfrak{g}_k$ (cf. [2, 3.3]). Note that the center of $\mathfrak{g}_k$ is contained in $\mathfrak{h}_k$ which is nontrivial only in the exceptional case, we have a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ arising from $\mathfrak{h}_\mathbb{C}$, which is equal to $\mathfrak{h}_k$ for non-exceptional cases, and isomorphic to $\mathfrak{h}_k/Z(\mathfrak{g}_k)$ for the exceptional case. Here $Z(\mathfrak{g}_k)$ means the center of $\mathfrak{g}_k$. Thus, we have a Cartan-decomposition relative to $\mathfrak{h}$: $\mathfrak{g} = \mathfrak{h} + \sum_{\alpha \in \Phi(\mathfrak{h})} \mathfrak{g}_\alpha$. Let $\Pi = \{\alpha_1, \dots, \alpha_r\}$ be a set of simple roots in the root system $\Phi := \Phi(\mathfrak{h})$. Then we have a Chevalley basis in the sense of the above meaning: $\{h_i, e_\alpha \mid 1 \leq i \leq r', \alpha \in \Phi\}$ where $r' = r - 1$ if $\mathfrak{g}$ is of the exceptional case, $r' = r$ if otherwise. Here e_α may be naturally regarded as $X_\alpha \otimes 1$ whether or not it's the exceptional case since the possible non-trivial center is in $\mathfrak{h}_k$. Additionally, each e_α defines a (nilpotent) linear transformation ade_α on $\mathfrak{g}$ by sending $y \in \mathfrak{g}$ to $[e_\alpha y]$. Thus, for any $a \in k$, set $x_\alpha(a) := \exp(\text{adae}_\alpha)$, which is easily seen to be a linear automorphism on $\mathfrak{g}$. Denote by $G(\mathfrak{h})$ the group generated by all $x_\alpha(a)$ which is called the Chevalley group of $\mathfrak{g}$ (constructed relative to $\mathfrak{h}$). Then $G(\mathfrak{h})$ coincides with G , which is independent of the choice of $\mathfrak{h}$ as k is algebraically closed ([23, III.4.1, IV. 1.1 and IV.1.2]). Actually, all Cartan subalgebras are conjugate under G ([23, IV.1.2]). Furthermore, $\text{Aut}(\mathfrak{g}) = G \rtimes \Gamma$ is a semi-direct product of G and the graph automorphism of Dynkin diagram of $\mathfrak{g}$ (cf. [23, III.Theorem 5.1 and §3]).

Let F_0 be a *standard* (or in other words, *splitting*) Frobenius-Lie morphism on $\mathfrak{g}$ which fixes the Chevalley basis of $\mathfrak{g}$. Then $\mathfrak{g}_0 = \mathfrak{g}^{F_0}$ is a *split* classical Lie algebra over $\mathbb{F}_q$ (of the same type as $\mathfrak{g}$) with respect to the Cartan subalgebra $\mathfrak{h}$. (Thus, $\mathfrak{g}_0$ has a Cartan decomposition of the form $\mathfrak{g}_0 = \mathfrak{h}^{F_0} + \sum_\alpha \mathfrak{g}_\alpha^{F_0}$). By Lang-Steinberg Theorem, it's proved in [10] that any Frobenius morphism F on $\mathfrak{g}$ is a split Frobenius morphism followed by a graph automorphism: $F = \gamma \cdot F'_0$ where $\gamma \in \Gamma$, $F'_0 \in G * F_0 := \{g * F_0 := g^{-1}F_0g \mid g \in G\}$. Moreover, such a decomposition is unique, and there is a one-to-one correspondence between the equivalence classes of Frobenius morphisms on $\mathfrak{g}$ and the conjugacy classes in Γ of graph automorphisms on $\mathfrak{g}$.

Remark 5.1. Precisely speaking, the number of (non-equivalent) Frobenius-Lie morphisms of a classical simple Lie algebra $\mathfrak{g}$ over k with $p > 3$ is as follows: one for types $A_1, B, C, G_2, F_4, E_7, E_8$; two for types $A_n (n > 1), D_n (n \geq 5), E_6$, and three for type D_4 . Especially, the representatives of equivalent Frobenius-Lie morphisms arises from $\sigma_0 = \text{Id}$, $\sigma_1 = (34)$ and $\sigma_2 = (13)$ where (ij) means the graph automorphism exchanging the i -th and j -th nodes in the standard Dynkin diagram of type D_4 , fixing the other nodes.

5.2. G -equivariant bijection on nilpotent orbits. We always fix a Frobenius-Lie morphism F on $\mathfrak{g}$ in the sequel, which has form $F = \gamma \cdot F'_0$. In the following we will discuss the relations between both of $\mathfrak{g}$ and $\mathfrak{g}_k$ in the nilpotent orbits and the Frobenius morphisms. First, we recall that G is naturally isomorphic to $\bar{G}/Z(\bar{G})$ where $Z(\bar{G})$ is the center of $\bar{G}$ which is in maximal tori of $\bar{G}$. Note that $Z(G) = \ker(\text{Ad})$ (cf. [3, 3.15]), and that $\mathfrak{g} = \mathfrak{g}_k/Z(\mathfrak{g}_k)$ with $Z(\mathfrak{g}_k) \subset \mathfrak{h}_k$. The following lemma is clear, which implies that we can replace G by $\bar{G}$ without affecting the structure of nilpotent orbit.

Lemma 5.2. *There is a G -equivariant bijection between $\mathcal{N}(\mathfrak{g})$ and $\mathcal{N}(\mathfrak{g}_k)$.*

Furthermore, such a bijection gives rise to a bijection between the nilpotent orbits of $\mathfrak{g}$ under G and the nilpotent orbits of $\mathfrak{g}$ under $\bar{G}$. Denote such a bijection by Ξ .

Now we look into the change of Frobenius morphisms from $\mathfrak{g}$ to $\mathfrak{g}_k$. In non-exceptional cases, there is no more words to say about this since $\mathfrak{g} = \mathfrak{g}_k$. Assume that $\mathfrak{g}$ is of type A_n with $p \mid n + 1$, then $\mathfrak{g} = \mathfrak{g}_k/Z(\mathfrak{g}_k)$. Let F be an arbitrary given Frobenius-Lie morphism on $\mathfrak{g}$. By the argument in §5.1, we may suppose that $F = \sigma \circ F'_0$, for $F'_0 = gF_0g^{-1}$ with $g \in G$. The Frobenius-Lie morphism F gives rise to a unique one on $\mathfrak{g}_k$ which is of form $\sigma \circ (\bar{g}\bar{F}_0\bar{g}^{-1})$, where $\bar{g}$ means a representative in $\bar{G}$ of g in connection with the quotient $\bar{G}/Z(\bar{G})$, and $\bar{F}_0$ means the standard Frobenius-Lie morphism on $\mathfrak{g}_k$ with respect to the Chevalley basis $\mathcal{B}_k$ of $\mathfrak{g}_k$. We denote by $\bar{F}$ such a Frobenius-Lie morphism

of $\mathfrak{g}_k$, arising from F . Obviously, the form of $\bar{F}$ is independent of the choice of $\bar{g} \in \bar{G}$ because $Z(G) = \ker(\text{Ad})$. We have the following

Lemma 5.3. *The correspondence $F \rightarrow \bar{F}$ is a bijection from the set of Frobenius-Lie morphisms on $\mathfrak{g}$ to the set of the Frobenius-Lie morphisms on $\mathfrak{g}_k$ of the form $\sigma \cdot (h\bar{F}_0h^{-1})$, $\sigma \in \Gamma$ and $h \in \bar{G}$.*

Combining the above lemmas, we have the following proposition which shows that the F -stability of nilpotent orbits of $\mathfrak{g}$ under G is equivalent to the $\bar{F}$ -stability of nilpotent orbits of $\mathfrak{g}_k$ under $\bar{G}$.

Proposition 5.4. *Let $\mathcal{O}$ be a nilpotent orbit in $\mathfrak{g}$, and F a Frobenius-Lie morphism on $\mathfrak{g}$. Then $\mathcal{O}$ is F -stable if and only if $\Xi(\mathcal{O})$ is $\bar{F}$ -stable.*

5.3. Weighted Dynkin diagrams and F -stability of nilpotent orbits. From this subsection on, the characteristic p of field k is assumed to be a good prime greater than 3 for $\mathfrak{g}$, which means that p is bigger than any coefficient of any root in Φ^+ expressed as a linear combination of simple roots. we will determine all F -stable nilpotent orbits in $\mathfrak{g}$. According to the previous arguments, we may replace $\mathfrak{g}$ by $\mathfrak{g}_k$.

Suppose G is a simply connected simple algebraic group. Then $\mathfrak{g}_k = \text{Lie}(G)$, where $\mathfrak{g}_k$ is defined as in the previous subsection. Let $\mathcal{O}_X$ be an arbitrary given nilpotent orbit in $\mathfrak{g}_k$. By Lemma 5.4, we only need to investigate the $\bar{F}$ -stability of $\mathcal{O}_X$ in $\mathfrak{g}_k$ under G , where the form of $\bar{F}$ is as defined as in Lemma 5.4.

We may attach its weighted Dynkin diagram, obtained by choosing a standard triple $\{Y, H, X\}$ (Jacobson-Morozov theorem, referred to [22], or [5] under stronger condition on p), a Cartan subalgebra $\mathfrak{h}$ containing H , and a set of simple roots Π making H dominant with $\langle \alpha_i, H \rangle \in \{0, 1, 2\}$ (referred to [21] or [20, §3.4] and [5, §5.6] under stronger condition on p ; [6] in characteristic 0), and then labelling every vertex corresponding to a simple root α with the integer $\langle \alpha, H \rangle$. This diagram depends only on the orbit $\mathcal{O}_X$. Observe that all Cartan subalgebras in $\mathfrak{g}_k$ are conjugate under adjoint action of G , and for the fixed T and then $\mathfrak{h}_k$, the corresponding root system $\Phi(\mathfrak{h}_k)$, the sets of simple roots are transitively acted by the Weyl group $W = N_G(T)/T$. Hence, without loss of generality, we may as well suppose that the weighted Dynkin diagram $\Delta(\mathcal{O}_X)$ are attached with an integer $\langle \alpha_i, H \rangle$ to the i th-node. The following Dynkin-Kostant theorem was proved in [21, 2.4] for a good characteristic p , in [5] and [20] for a stronger characteristic p , in [6] for characteristic 0.

Theorem 5.5. (Kostant) *Let $\mathcal{O}_1$ and $\mathcal{O}_2$ be nilpotent orbits in $\mathfrak{g}_k$. Then $\mathcal{O}_1 = \mathcal{O}_2$ if and only if $\Delta(\mathcal{O}_1) = \Delta(\mathcal{O}_2)$.*

First, we investigate the situation when $\bar{F}$ is a standard Frobenius-Lie morphism $\bar{F}_0$.

Lemma 5.6. (a) *All nilpotent orbits $\mathcal{O}_X$ in $\mathfrak{g}_k$ are $\bar{F}_0$ -stable.*

(b) *The nilpotent orbit $\mathcal{O}_X$ under G is $\bar{F}$ -stable if and only if it's γ -stable.*


Proof. (a) Suppose $\{X, H, Y\}$ are a standard triple in sense of Jacobson-Morozov theorem (cf. [22] and [5]). As discussed in the previous paragraph, we might as well suppose that the weighted Dynkin diagram $\Delta(\mathcal{O}_X)$ admits the number $\langle \alpha_i, H \rangle$ to the i -th node for all i . Hence we may write $H = \sum_i a_i h_{\alpha_i}$ with all $a_i \in \mathbb{Z}_p$. Hence $\bar{F}_0(H) = H$. Observe that $\bar{F}_0$ is an $\mathbb{F}_q$ -Lie isomorphism, thereby $\{\bar{F}_0(X), \bar{F}_0(H), \bar{F}_0(Y)\}$ is still a standard triple. Hence $\Delta(\mathcal{O}_X) = \Delta(\mathcal{O}_{\bar{F}_0(X)})$. By Dynkin-Kostant theorem, we know $\bar{F}(\mathcal{O}_X) = \mathcal{O}_{\bar{F}_0(X)} = \mathcal{O}_X$.

(b) Recall that $\bar{F}'_0(\mathcal{O}_X) = g^{-1} \circ \bar{F}_0 \circ g(G.X)$. By Statement (a), $\mathcal{O}_X$ is a $\bar{F}'_0$ -stable. $\square$

Next, we will suppose that $\bar{F}$ is a twist Frobenius-Lie morphism, i.e. $\bar{F} = \gamma \cdot \bar{F}'_0$ for a non-trivial graph automorphism. Surely, this happens only in the case of types A_n, D_n and E_6 . Recall that for $\mathfrak{g}_k$ of type A_n and D_n , each nilpotent orbit uniquely corresponds to a partition of n and $2n$


respectively. Precisely speaking, nilpotent orbits in $\mathfrak{sl}_n$ are in one-to-one correspondence with the set $\mathcal{P}(n)$ of partitions of n . Nilpotent orbits in $\mathfrak{so}_{2n}$ are parameterized by partitions of $2n$ in which even parts occur with even multiplicity, except that "very even" partitions $\mathbf{d} := [d_1, d_2, \dots, d_{2n}]$ (those with only even parts, each having even multiplicity) correspond to two orbits, denoted by $\mathcal{O}_{[d_1, d_2, \dots, d_{2n}]}^I$ and $\mathcal{O}_{[d_1, d_2, \dots, d_{2n}]}^{II}$. For their weighted Dynkin diagrams, we have the following results.

Proposition 5.7. ([20, §3.4 and §5.7] and [6]) (1) For $\mathfrak{g}_k = \mathfrak{sl}_n$, let $\mathcal{O} = \mathcal{O}_{[d_1, \dots, d_n]}$. Then the weighted Dynkin diagram $\Delta(\mathcal{O})$ is given by




where h_i has the following meaning. Replace each part d_i by the list $d_i - 1, d_i - 3, \dots, -d_i + 1$ of weights in the $\mathfrak{sl}_2$ -module of dimension d_i and combine the resulting lists into one, rearranging its terms in non-increasing order as $h_1, \dots, h_n$.

(2) For $\mathfrak{g}_k = \mathfrak{so}_{2n}$, $n > 4$, let $\mathcal{O} = \mathcal{O}_{\mathbf{d}}$ for $\mathbf{d} = [d_1, \dots, d_{2n}]$. If $\mathbf{d}$ is not very even, then the weighted Dynkin diagram of $\mathcal{O}_{\mathbf{d}}$ is




If $\mathbf{d}$ is very even, then the weighted diagrams of $\mathcal{O}_{\mathbf{d}}^I$ and $\mathcal{O}_{\mathbf{d}}^{II}$ are given by




Here the meaning of h_i is as in the case $\mathfrak{sl}_n$.

(3) Let $\mathfrak{g}_k = \mathfrak{so}_8$. We have the following complete list of weighted Dynkin diagrams of all nilpotent orbits.


(i) The weighted Dynkin diagrams corresponding to the nilpotent orbits with partitions $\mathbf{d} = [7, 1], [5, 3], [2^2, 1^4]$ and $[1^8]$ are listed respectively as follows:



(ii) The weighted Dynkin Diagrams corresponding to the nilpotent orbits $\mathcal{O}_{[4^2]}^I$, $\mathcal{O}_{[4^2]}^{II}$ and $\mathcal{O}_{[5, 1^3]}$ are listed respectively in the following



(iii) The weighted Dynkin Diagrams corresponding to the nilpotent orbits $\mathcal{O}_{[2^4]}^I, \mathcal{O}_{[2^4]}^{II}$ and $\mathcal{O}_{[3,1^5]}$ are listed respectively in the following



(4) When $\mathfrak{g}_k$ is of type E_6 , there are 21 nilpotent orbits whose weighted Dynkin diagrams admit symmetric labels in accordance with the symmetry of the Dynkin diagram (to see [20, Page 202]).

We have the following result.

Theorem 5.8. *Let G be a simply connected simple algebraic group over k with a good characteristic $p > 3$ associated to the root system of G , $\mathfrak{g}_k$ the Lie algebra of G . Assume a Frobenius-Lie morphism $\bar{F} = \sigma \circ (\bar{F}'_0)$ with $\bar{F}'_0 = g \cdot F_0 g^{-1}$ as defined as in the previous subsection. For $\mathcal{O}_X$ a non-zero nilpotent orbit in $\mathfrak{g}_k$ we have*

(1) *If $\mathfrak{g}_k$ is of type A_n or type E_6 , then $\mathcal{O}_X$ are $\bar{F}$ -stable.*

(2) *Let $\mathfrak{g}_k$ is of type D_n with $n > 4$. Then $\mathcal{O}_X$ is $\bar{F}$ -stable for all such $\bar{F}$ with exception when $\mathcal{O}_X$ is very-even, and $\bar{F}$ is twist. In the exceptional cases, the nilpotent orbits $\mathcal{O}_{\mathbf{d}}^I$ and $\mathcal{O}_{\mathbf{d}}^{II}$ are exchanged under a twist $\bar{F}$.*

(3) *Let $\mathfrak{g}_k$ is of type D_4 . Keep the notations as in Remark 5.1,*

(i) *$\bar{F}_0$ stabilizes all nilpotent orbits.*

(ii) *$\bar{F}_1 : \mathcal{O}_{[4^2]}^I \longleftrightarrow \mathcal{O}_{[4^2]}^{II}$, and $\mathcal{O}_{[2^4]}^I \longleftrightarrow \mathcal{O}_{[2^4]}^{II}$, keeping the other nilpotent orbits fixed.*

(iii) *$\bar{F}_2 : \mathcal{O}_{[2^4]}^{II} \longleftrightarrow \mathcal{O}_{[3,1^5]}$, and $\mathcal{O}_{[4^2]}^{II} \longleftrightarrow \mathcal{O}_{[5,1^3]}$, keeping the other nilpotent orbits fixed.*

Proof. By Lemma 5.6, we only need to work with a twist Frobenius-Lie morphism $\bar{F}$. By Lemma 5.6 again, the $\bar{F}$ -stability of $\mathcal{O}_X$ is equivalent to its γ -stability. As having discussed above, we might as well suppose that the weighted Dynkin diagram arises from a standard triple $\{X, H, Y\}$ and the fixed Cartan subalgebra $\mathfrak{h}_k$ along with a set of simple root $\Pi(\mathfrak{h}_k)$ in the root system $\Phi(\mathfrak{h}_k)$ such that $\langle \alpha_i, H \rangle \in \{0, 1, 2\}$. Note that $\mathfrak{g}_k$ has a Chevalley basis, a perfect analogue of a Chevalley basis of $g_{\mathbb{C}}$. Naturally, the graph automorphism γ of the Dynkin diagram satisfies:

$$\langle \alpha_{\gamma(i)}, \gamma(H) \rangle = \langle \alpha_i, H \rangle.$$

Observe that the Lie automorphism γ implies that $\{\gamma(X), \gamma(H), \gamma(Y)\}$ is also a standard triple. In the meantime, $\gamma(\Pi)$ is a set of simple roots of $\Phi(\mathfrak{h}_k)$. Hence $\langle \alpha_{\gamma(i)}, \gamma(H) \rangle$ gives rise the weighted Dynkin diagram of the nilpotent orbit of $\gamma(X)$. Comparing the weighted Dynkin diagrams for type A_n , D_n and E_6 (cf. [20, 3.4 and 5.7] and [6, 3.6, 5.3 and 8.4]), we know that $\Delta(X)$ coincides with $\Delta(\gamma(X))$ for all cases except that $\mathcal{O}_X$ is very even with a very even partition $\mathbf{d}$ which happens in the case of D_n . In the exceptional cases, $\bar{F}(\Delta(\mathcal{O}_{\mathbf{d}}^I)) = \Delta(\mathcal{O}_{\mathbf{d}}^{II})$ for D_n with $n > 4$. When $\mathfrak{g}_k$ is of D_4 , then it's not hard to get the result by comparing the weighted Dynkin diagrams in Proposition 5.7. $\square$

Remark 5.9. Denote $\bar{F}_3$ by the Frobenius map arising from the graph automorphism (14). Then $\bar{F}_3$ is equivalent to $\bar{F}_2$. And the effect of $\bar{F}_3$ -action makes exchanges: $\mathcal{O}_{[24]}^I \longleftrightarrow \mathcal{O}_{[3,1^5]}$, and $\mathcal{O}_{[4^2]}^I \longleftrightarrow \mathcal{O}_{[5,1^3]}$, keeping the other nilpotent orbits fixed.

By Lemma 5.3 and Proposition 5.4, we have the following theorem.

Theorem 5.10. *Let $\mathfrak{g}$ be a classical simple Lie algebra over $k = \bar{\mathbb{F}}_q$, defined over a finite field $\mathbb{F}_q$ of a good characteristic $p > 3$, with correspondence to Frobenius map F , let G the adjoint group of $\mathfrak{g}$. Then the following statements hold:*

(1) $F = \sigma \cdot (gF_0g^{-1})$, where $\sigma \in \Gamma$, $g \in G$ and F_0 is a standard Frobenius morphism associated with a Chevalley basis (in the sense of §5.1).

(2) For a nilpotent orbit $\mathcal{O}_X$ in $\mathfrak{g}$ under the action of G , all statements in Theorem 5.8 hold.

6. $\mathbb{F}_q$ -RATIONAL STRUCTURE ON NILPOTENT ORBITS

Let G be a connected semi-simple algebraic group over an algebraically closed field F of characteristic $p > 3$, which is defined over $\mathbb{F}_q$ with corresponding Frobenius map F . Let $F_{\mathfrak{g}}$ be the corresponding Frobenius-Lie morphism of $\mathfrak{g} = \text{Lie}(G)$. We have

Lemma 6.1. *For any $g \in G$ and $X \in \mathfrak{g}$, $F_{\mathfrak{g}}(g \cdot X) = F(g) \cdot F_{\mathfrak{g}}(X)$.*

Proof. Let T be a fixed maximal torus of G , and Δ be the rt system of G associated with T . We have the corresponding complex semisimple Lie algebra $\mathfrak{g}_{\mathbb{C}}$ with triangular decomposition $\mathfrak{g}_{\mathbb{C}} = \mathfrak{n}_{\mathbb{C}}^- + \mathfrak{h}_{\mathbb{C}} + \mathfrak{n}_{\mathbb{C}}^+$. Take in $\mathfrak{g}_{\mathbb{C}}$ a Chevalley basis $\{X_{\alpha}, X_{-\alpha}, H_i \mid \alpha \in \Delta_+, i = 1, 2, \dots, l := \dim(T)\}$. By [25], there is a faithful finite-dimensional representation (π, V) of $\mathfrak{g}_{\mathbb{C}}$ with the weight lattice equal to the character group $X(T)$, and an admissible $\mathbb{Z}$ -lattice M of $\mathfrak{g}_{\mathbb{C}}$ -module V . Then G can be realized as the Chevalley group G_V which is a closed subgroup of $GL(V_k)$ for $V_k = M \otimes k$. Recall that G_V is generated by elements: $x_{\alpha}^{\pi}(t) = \sum_{n=0}^{\infty} \pi(X_{\alpha}^{(n)})t^n$, for $\alpha \in \Delta$ and $t \in k$. Here $X_{\alpha}^{(n)} = \frac{X_{\alpha}^n}{n!}$. So we only need to prove that $F_{\mathfrak{g}}(g \cdot X) = F(g) \cdot F_{\mathfrak{g}}(X)$ for $g = x_{\alpha}^{\pi}(t)$ and $X = X_{\beta}$ or H_i for all $\alpha, \beta \in \Delta$, $i = 1, \dots, l$ and $t \in k$. Notice that F is decided by an element of $\text{Aut}(\Gamma)$ ([5, 1.19]), say σ , where Γ is the Dynkin diagram of Δ . So we have $F(x_{\alpha}^{\pi}(t) = x_{\sigma(\alpha)}^{\pi}(t^q)$. In the same time, $F_{\mathfrak{g}} = \sigma \cdot F_0$, where F_0 is the splitting Frobenius-Lie morphism associated with $\mathfrak{h} = \text{Lie}(T)$. We identify $\mathfrak{g}$ with $\text{Lie}(G_V)$, which is a linear Lie subalgebra of $\mathfrak{gl}(V_k)$. By [2], $\mathfrak{g} = \mathfrak{g}_{\pi} \otimes_{\mathbb{Z}} k$, where $\mathfrak{g}_{\pi} = \sum \mathbb{Z}X_{\alpha} + \mathfrak{h}_{\pi}$, and $\mathfrak{h}_{\pi} = \{h \in \mathfrak{h}_{\mathbb{C}} \mid \alpha(h) \in \mathbb{Z}, \forall \alpha \in X(T)\}$. So we can suppose the splitting Frobenius mapping F_0 is associated with a Chevalley basis $\mathcal{B}_{\pi} = \{X_{\alpha}, H'_i \mid \alpha \in \Delta, i = 1, \dots, l\}$. Then $F_{\mathfrak{g}}(\sum_{\alpha} a_{\alpha}X_{\alpha} + \sum_i b_i H'_i) = \sum_{\alpha} a_{\alpha}^q X_{\sigma(\alpha)} + \sum_i b_i^q H'_i$ for all $a_{\alpha}, b_i \in k$.

Then we have $x_{\alpha}^{\pi}(t) \cdot Y = \sum_{n=0}^{\infty} \text{ad}(X_{\alpha}^{(n)})(Y)$ for $Y = X_{\beta}$ or H'_i .

Case 1. $Y = X_{\beta}$.

Observe that the lengths of root chains in Δ are 4 at most, which is less than p . And furthermore, $\text{ad}(X_{\alpha})^n(X_{\beta}) = C_{\alpha, \beta, n} X_{n\alpha + \beta}$ if $n\alpha + \beta \in \Delta$, and 0 if $n\alpha + \beta$ is not a root, with $C_{\alpha, \beta, n} \in \mathbb{Z}$ satisfying that

$$(6.1.1) \quad \text{ad}(X_{\sigma(\alpha)})^n(X_{\sigma(\beta)}) = C_{\alpha, \beta, n} X_{n\sigma(\alpha) + \sigma(\beta)} \text{ if } n\alpha + \beta \in \Delta, \text{ and } 0 \text{ if } n\alpha + \beta \text{ is not a root.}$$

Thus, we have

$$(6.1.2) \quad \begin{aligned} & F_{\mathfrak{g}}\left(\frac{\text{ad}(X_{\alpha}^n)(X_{\beta})}{n!}\right) \\ &= C_{\alpha, \beta, n} \frac{X_{n\sigma(\alpha) + \sigma(\beta)}}{n!} \\ &= \frac{\text{ad}(X_{\sigma(\alpha)})^n(X_{\sigma(\beta)})}{n!}. \end{aligned}$$

Hence we have

$$\begin{aligned}
 & F_G(x_\alpha^\pi(t) \cdot F_{\mathfrak{g}}(X_\beta)) \\
 &= x_{\sigma(\alpha)}^\pi(t^q) \cdot X_{\sigma(\beta)} \\
 (6.1.3) \quad &= \sum_{n=0}^{\infty} \frac{\text{ad}(X_{\sigma(\alpha)})^n(X_{\sigma(\beta)})}{n!} t^{nq}
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 & F_{\mathfrak{g}}(x_\alpha^\pi(t) \cdot X_\beta) \\
 &= F_{\mathfrak{g}}\left(\sum_{n=0}^{\infty} \frac{(\text{ad}(X_\alpha)^n(X_\beta))}{n!} t^n\right) \\
 (6.1.4) \quad &= \sum_{n=0}^{\infty} F_{\mathfrak{g}}\left(\frac{(\text{ad}(X_\alpha)^n(X_\beta))}{n!}\right) t^{nq}.
 \end{aligned}$$

By (6.1.2), we have $F_{\mathfrak{g}}(g \cdot Y) = F(g)F_{\mathfrak{g}}(Y)$ for $g = x_\alpha^\pi(t)$ and $Y = X_\beta$.

Case 2. $Y = H'_i$.

In this case, $\text{ad}(X_\alpha)H'_i = -\langle \alpha, \alpha_i^\vee \rangle X_\alpha$ and $\text{ad}(X_\alpha)^n H'_i = 0$ for $n > 1$. So it's more easily seen by the same argument that $F_{\mathfrak{g}}(g \cdot Y) = F(g) \cdot F_{\mathfrak{g}}(Y)$ for $g = x_\alpha^\pi(t)$ and $Y = H'_i$. $\square$

Thus combining the above lemma and Lemma 1.7, we have

Proposition 6.2. *Let G be a connected semi-simple algebraic group over an algebraically closed field F of characteristic $p > 3$, which is defined over $\mathbb{F}_q$ with corresponding Frobenius map F . Let $F_{\mathfrak{g}}$ be the corresponding Frobenius-Lie morphism of $\mathfrak{g} = \text{Lie}(G)$. Then any $F_{\mathfrak{g}}$ -stable orbit in $\mathfrak{g}$ admits $\mathbb{F}_q$ -rational structure, thereby admits $F_{\mathfrak{g}}$ -fixed points. Especially, if $\bar{G}$ is a simply-connected simple algebraic group, $\mathfrak{g}_k$ is the Lie algebra of G , then $\bar{F}$ as defined as in Lemma 5.3 is a Frobenius-Lie morphism arising from a Frobenius map on G . Therefore, any $\bar{F}$ -stable orbit in $\mathfrak{g}_k$ under the action of $\bar{G}$ admits an $\mathbb{F}_q$ -rational structure, thereby admits $\bar{F}$ -fixed points.*

Furthermore, we have the following:

Proposition 6.3. *Let $\mathfrak{g}$ be a classical simple Lie algebra over $k = \bar{\mathbb{F}}_q$, defined over a finite field $\mathbb{F}_q$ of a good characteristic $p > 3$, with correspondence to Frobenius map F , let G the adjoint group of $\mathfrak{g}$. For an F -stable nilpotent orbit $\mathcal{O}$ which is listed in 5.8, $\mathcal{O}$ admits an $\mathbb{F}_q$ -rational structure, thereby admits F -fixed points.*

Proof. By the structure of G (cf. §5.1), we may define a Frobenius map F_G on G such that the Frobenius-Lie morphism on $\mathfrak{g}$ arising from F_G is just F . Observe that all Cartan subalgebra are conjugate under G ([23, IV.1.2]). We might as well assume that $\mathfrak{h}$ is such a Cartan subalgebra that $F = \sigma \circ F_0$ for a splitting Frobenius-Lie morphism F_0 , associated with $\mathfrak{h}$. As stated in §5.1, there are generators in G : $x_\alpha(a) = \exp(\text{ada}_{e_\alpha})$ for $a \in k$. We may define F_G via $F_G(x_\alpha(a)) = x_{\sigma(\alpha)}(a^q)$. Then for every orbit in $\mathfrak{g}$ under the action of G , the Frobenius maps are admissible with the action of G . This is to say, $F(g.X) = F_G(g).F(X)$ because F is a Frobenius-Lie morphism. $\square$

In the following theorem, we continue to assume:

- (1) $\mathfrak{g}$ is a classical simple Lie algebra, and G is an adjoint group of $\mathfrak{g}$. F is an arbitrary given Frobenius-Lie morphism on $\mathfrak{g}$.
- (2) $\bar{G}$ is a simply connected algebraic group and $\mathfrak{g}_k = \text{Lie}(G)$. $\bar{F}$ is as defined as in Lemma 5.8.

Summing up the above results, we can state the following theorem.

Theorem 6.4. *Let $\mathcal{O}_X$ is a nilpotent orbit in $\mathfrak{g}$ under G -action, or in $\mathfrak{g}_k$ under $\bar{G}$ -action over $k = \bar{\mathbb{F}}_q$, defined over a finite field $\mathbb{F}_q$ of a good characteristic $p > 3$. Then $\mathcal{O}_X$ admits $\mathbb{F}_q$ -rational structure if and only if $\mathcal{O}_X$ is F -stable (resp. $\bar{F}$ -stable for $\mathfrak{g}_k$), which is completely listed in Theorem 5.8.*

REFERENCES

- [1] Benson, D., *Modular representation theory: new trends and methods*, Springer-Verlag, LNM 1081, 1984.
- [2] Borel, A., *Properties and linear representations of Chevalley groups*, pages 1-55, in "Seminar on algebraic groups and related finite groups, Lecture Notes in Mathematics 131, Springer-Verlag, Berlin, 1970.
- [3] Borel, A., *Linear algebraic groups*, Springer-Verlag, 1991.
- [4] Carlson, J., *The variety of an indecomposable module is connected*, Inven. Math. (1987)
- [5] Carter, R.W., *Finite groups of Lie type*, Wiley-Interscience, 1985.
- [6] Collingwood, D.H. and McGovern, W.M., *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand Reinhold, New York, 1993.
- [7] Curtis, C.W. and Reiner, I., *Methods of representation theory. With applications to finite groups and orders*, Vol.I, A Wiley-Interscience Publication, New York, 1981.
- [8] Deng, B. and Du, J., *Frobenius morphisms and representations of algebras*, Trans. Amer.Math. Soc. (to appear).
- [9] Digne, F. and Michel, J., *Representations of finite groups of Lie type*, London Math. Soc., Student Texts, 21. Cambridge University Press, Cambridge, 1991.
- [10] Du, J. and Shu B., *On finite Lie algebras and their representations*, preprint.
- [11] Friedlander, E. M. and Parshall, B. J., *Cohomology of algebraic and related finite groups*, Inv. Math. 74 (1983), 85-117.
- [12] Friedlander, E. M. and Parshall, B. J., *Support varieties for restricted Lie algebras*, Inv. Math. 86 (1986), 553-562.
- [13] Friedlander, E. M. and Parshall, B. J., *Geometry of p -unipotent Lie algebras*, J. Algebra 109 (1987), 25-45.
- [14] Friedlander, E. M. and Parshall, B. J., *Modular representations theory of Lie algebras*, Amer. J. Math. 110 (1988), 117-126.
- [15] Geck, M., *An introduction to algebraic geometry and algebraic groups*, Oxford University Press, 2003.
- [16] Humphreys, J.E., *Algebraic groups and modular Lie algebras*, Memoirs of the American Mathematical Society, vol. 71, 1967.
- [17] Jantzen, J.C., *Representations of algebraic groups*, Academic Press, 1987.
- [18] Jantzen, J.C., *Restricted Lie algebra cohomology*, pp. 91-108, in *Algebraic groups: Proceedings of a conference in honor of T. A. Springer*, Lecture Notes in Mathematics 1271, Springer-Verlag, 1987.
- [19] Jantzen, J.C. *Nilpotent orbits in representation theory*, pages 1-221, in "Lie Theory, Lie algebras and representations", Progress in Mathematics 228, Birkhäuser, Boston.Basel.Berlin, 2004.
- [20] McGovern, W.M., *The adjoint representation and the adjoint action*, in *Algebraic Quotients. Torus actions and cohomology. The adjoint representation and the adjoint action*, Springer 2002, 150-238.
- [21] Premet, Alexander, *Nilpotent orbits in good characteristic and the Kempf-Rousseau theory*, Journal of Algebra, 260(2003), 338-366.
- [22] Premet, Alexander, *An analogue of the Jacobson-Morozov theorem for reductive groups of good characteristics*, Trans. Amer. Math. Soc. 347 (1995) 2961-2988.
- [23] Seligman, G.B., *Modular Lie algebras*, Springer-Verlag, Berlin Heidelberg New York, 1967.
- [24] Springer, T.A. and Steinberg, R., *Conjugacy class*, pages 167-266, in "Seminar on algebraic groups and related finite groups, Lecture Notes in Mathematics 131, Springer-Verlag, Berlin, 1970.
- [25] Steinberg, R., *Lectures on Chevalley groups*, Yale University, 1968.

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, ANKARA, TURKEY
E-mail address: semra@math.metu.edu.tr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904
E-mail address: bjp8w@virginia.edu

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200062, CHINA.
E-mail address: bshu@euler.math.ecnu.edu.cn